

AD-A064 302

FLORIDA STATE UNIV TALLAHASSEE DEPT OF OCEANOGRAPHY
RAY CURVATURE AND REFRACTION OF WAVE PACKETS.(U)
SEP 78 J E BREEDING

F/G 8/3

N00014-77-C-0329

UNCLASSIFIED

TR-JEB-3

NL

1 OF 1
AD
A0 64 302



AD A 064302

DDC FILE COPY

LEVEL 1

(8)
F


RAY CURVATURE AND REFRACTION OF WAVE PACKETS

by J. Ernest Breeding, Jr.

TECHNICAL REPORT NO. JEB-3

Department of Oceanography
Florida State University

September, 1978



Approved for Public Release;
Distribution Unlimited

Sponsored by the Geography Programs, Earth Sciences Division,
Office of Naval Research, under Contract No. N00014-77-C-0329

79 01 29 014


Technical Report No. JEB-3

Department of Oceanography
Florida State University

RAY CURVATURE AND REFRACTION
OF WAVE PACKETS

by

J. Ernest Breeding, Jr.
Department of Oceanography
Florida State University
Tallahassee, Florida 32306



September, 1978

This research was supported by the Geography Programs, Earth Sciences Division, Office of Naval Research under Contract No. N00014-77-C-0329. Reproduction in whole or in part is permitted for any purpose of the United States Government. Approved for public release; distribution unlimited.

79 01 29 014

ABSTRACT

An expression for the ray curvature of a wave packet is derived which is suitable for use in wave prediction programs. The ray curvature of the wave packet is found to vanish if the packet direction becomes either perpendicular or parallel to the wave speed contours, assuming the wavelet direction is not parallel to the contours. For parallel water depth contours, this means that as a hydron moves into shoaling water refraction tends eventually to turn the hydron so that it is directed either perpendicular or parallel to the shoreline. The first case is similar to monochromatic waves. It is the result for hydrons which begin in deep water if the angle of incidence is between 0° and 74.8° with respect to the contour normal. However, for deep water angles of incidence equal to or greater than 74.8° the hydrons are turned and move parallel to shore in water of intermediate depth. The packet ray curvature approaches infinity as the wavelet direction, but not the hydron direction, becomes parallel to the wave speed contours. The result is total reflection of the waves. Total reflection occurs if a hydron is moving into deeper water and its initial direction exceeds a critical angle. At the reflection point the hydron direction becomes perpendicular to the water depth contours. Further, the hydron velocity goes to zero, which is consistent with a particle concept. As in quantum mechanics, the wave-particle duality is encountered.

File	White Section
EDS	Self Section <input type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
SPECIAL	
A	

Acknowledgments. I thank Franklyn C. W. Olson, Kenneth C. Matson, and Kimberly Crane Oppenheimer for their help in preparing this paper. The research was supported by the Geography Programs, Earth Sciences Division, Office of Naval Research under Contract No. N00014-77-C-0329.

TABLE OF CONTENTS

	<u>Page</u>
Abstract	ii
Acknowledgments	iii
Table of Contents	iv
List of Figures	v
Introduction	1
Ray Curvature for Wave Packets	1
Packet Ray Curvature for Locally Parallel Wave Speed Contours	2
Properties of the Packet Ray Curvature and Snell's Law	3
Examples of Wave Packet Refraction-Hydrons	5
Waves from Deep Water.....	5
Reflection Points	7
Summary	10
References	13
Distribution List	14

LIST OF FIGURES

Page

Figure 1.	Hydron trajectories for a 20 sec wave period for waves beginning in deep water	6
Figure 2.	Monochromatic rays for comparison with the hydron trajectories in Figure 1.....	6
Figure 3.	The variation of the geometric group speed with kh for hydrons beginning in deep water	8
Figure 4.	Hydron trajectories for a 20 sec wave period for waves which begin at an intermediate water depth	9
Figure 5.	Variations of the wavelet direction γ , hydron direction θ , the speed $U = d\omega/dk$, and the geometric group speed $G = U \cos (\theta - \gamma)$ up to the reflection point for ray number 2 in Figure 4	11
Figure 6.	The ratio of the hydron ray curvature to its absolute initial value as a function of the wavelet direction up to the reflection point for ray number 2 in Figure 4	12

Introduction

The conventional definition of group speed U has been defined as

$$U = d\omega/dk \quad (1)$$

where ω is the angular frequency and k is the wave number. This equation defines the speed of the group in the direction of the wavelet velocity. The geometric group speed G was defined by Breeding (1978) as

$$G = U \cos \phi \quad (2)$$

where

$$\phi = \theta - \gamma \quad (3)$$

The direction of movement of the wave packet is represented by θ , and the direction the wavelets move within the packet is specified by γ .

Breeding (1978) has shown that wave packets refract according to Snell's law with the geometric group velocity. This refraction law determines the wave path of constructive interference. At each point of the wave packet trajectory the wavelet direction is determined by Snell's law with phase velocity.

In this paper an expression is derived for the ray curvature of a wave packet and its properties are described. Examples of gravity water waves will be presented to show how G and U differ and to demonstrate the important features of wave packet refraction. In addition, wave packet and monochromatic trajectories will be compared for waves which begin in deep water.

Ray Curvature for Wave Packets

The ray curvature κ_v of a ray moving with phase speed v was derived by Munk and Arthur (1952) and Arthur, et al (1952) as

$$\kappa_v = \frac{dy}{ds_v} = \frac{1}{v} \left(\sin \gamma \frac{\partial v}{\partial x} - \cos \gamma \frac{\partial v}{\partial y} \right) \quad (4)$$

where x, y are the Cartesian coordinates, γ is the direction of the ray with respect to the positive x -axis, and s_v is the arc length along the ray.

The ray curvature κ_G for the trajectory of a wave packet is given by

$$\kappa_G = \frac{d\theta}{ds_G} = \frac{1}{G} \left(\sin \theta \frac{\partial G}{\partial x} - \cos \theta \frac{\partial G}{\partial y} \right) \quad (5)$$

where θ is the direction of the packet ray with respect to the positive x-axis and s_G is the arc length along the ray.

Packet Ray Curvature for Locally Parallel Wave Speed Contours

In deriving an expression for the packet ray curvature the wave speed contours are assumed to be parallel. This assumption greatly simplifies the derivation. Further, it leads to an expression suitable for making calculations from which the important properties of the packet ray curvature can be determined. Accordingly, the xy-coordinate system is chosen such that the y-derivatives of v , U , and G are zero. The space derivative of G is then

$$\frac{dG}{dx} = \frac{dU}{dx} \cos \phi - U \sin \phi \left(\frac{d\theta}{dx} - \frac{d\gamma}{dx} \right) \quad (6)$$

Expressions for the space derivatives of θ and γ are obtained from (4) and (5)

where

$$dx = ds_v \cos \gamma \quad (7)$$

$$dx = ds_G \cos \theta \quad (8)$$

$$\text{Thus} \quad d\gamma/dx = [(dv/dx)/v] \tan \gamma \quad (9)$$

$$d\theta/dx = [(dG/dx)/G] \tan \theta \quad (10)$$

After (9), (10), and (6) are substituted into (5) and the result is simplified, the packet ray curvature is found to be

$$\kappa_G = \frac{[(dU/dx)/U] + [(dv/dx)/v] \tan \phi \tan \gamma}{\csc \theta + \tan \phi \sec \theta} \quad (11)$$

This expression for the packet ray curvature can be used in wave prediction programs even if the wave speed contours are not everywhere parallel. From a practical viewpoint, it is only necessary that the wave speed contours be locally parallel in the vicinity of each ray point. Eq. (11) is used by Breeding et al 1978

in computing wave packet trajectories of gravity water waves. To use (11) the coordinate system is rotated at each ray point so that the y-axis is parallel with the water depth contours and the positive x-axis is directed toward deeper water.

Properties of the Packet Ray Curvature and Snell's Law

The ray curvature of a wave packet defined by (11) exhibits some very remarkable properties. It is assumed that v , U , and their derivatives are continuous and finite. However, under various conditions the trigonometric terms of the equation can become infinite or have indeterminate forms. The value of κ_G approaches zero as the wave packet direction θ becomes either parallel or perpendicular to the wave speed contours, provided the wavelet direction γ is not parallel to the contours. This means that given a sufficiently long path, refraction tends to turn the wave packet so that it is directed either parallel or perpendicular to the wave speed contours. If θ is neither parallel nor perpendicular to the wave speed contours, then κ_G approaches infinity as γ becomes parallel to the wave speed contours. In this case, due to the value of γ , the wave packet undergoes total reflection.

To determine the value of κ_G when there are indeterminate forms it is necessary to consider the variations of θ and γ as the indeterminate forms are approached. For example, (11) contains an indeterminate form when θ becomes perpendicular to the wave speed contours while γ becomes parallel to the contours. If γ approaches parallelism to the contours faster than θ approaches the perpendicular to the contours the value of κ_G becomes infinite.

The relationship between θ and γ due to refraction is clearly seen by integrating the ray curvature expressions (4) and (5). The y-derivatives being zero, integration of (5) leads to

$$(\sin \theta) / [U \cos (\theta - \gamma)] = C \quad (12)$$

which is Snell's law for a wave packet where C is a constant. Snell's law with

phase velocity, which determines γ , is obtained by integrating (4). The cosine term in (12) can be replaced by the identity for the difference of two angles and the terms rearranged to yield

$$\tan \theta = (UC \cos \gamma) / (1 - UC \sin \gamma) \quad (13)$$

where the variation of θ appears only on the left side of the equation.

It is interesting to note that θ becomes zero if $\gamma = (2m + 1) (\pi/2)$ where m is an integer. Thus if the wavelet direction becomes parallel to the wave speed contours the wave packet direction becomes perpendicular to the contours. Further, note that $\theta = (2m + 1) (\pi/2)$ if $UC \sin \gamma = 1$. For this case the wave packet direction is parallel to the wave speed contours.

Snell's law can be used to derive an expression for $\cos (\theta - \gamma)$. Eq. (13) is substituted into the identity for $\tan (\theta - \gamma)$ and the result is simplified to obtain

$$\tan (\theta - \gamma) = (UC - \sin \gamma) / \cos \gamma \quad (14)$$

In terms of initial values, Snell's law with phase velocity can be written

$$\sin \gamma = v_r \sin \gamma_i \quad (15)$$

where $v_r = (v/v_i)$ and the subscript i denotes an initial value. Before refraction it is assumed that $\theta_i = \gamma_i$. Then

$$C = (\sin \gamma_i) / U_i \quad (16)$$

When (15) and (16) are substituted into (14), it is found that

$$\tan (\theta - \gamma) = \frac{(U_r - v_r) \sin \gamma_i}{(1 - v_r^2 \sin^2 \gamma_i)^{1/2}} \quad (17)$$

where $U_r = (U/U_i)$. This result can be transformed by the use of an identity to

$$\cos (\theta - \gamma) = \left[1 + \frac{(U_r - v_r)^2 \sin^2 \gamma_i}{1 - v_r^2 \sin^2 \gamma_i} \right]^{-1/2} \quad (18)$$

Eqs. (1), (2), and (18) provide a useful means of computing the geometric group speed.

Examples of Wave Packet Refraction - Hydrons

To demonstrate the properties of wave packet refraction, examples of gravity water waves will be considered. Gravity water waves are particularly suited as examples because of their highly dispersive nature. The term 'hydron,' suggested by Purser and Synge (1962) and Synge (1962), will be used to denote the wave packet of water waves.

Waves from Deep Water

In Figure 1 hydron trajectories are shown for waves beginning in deep water (water depth greater than one half the wavelength). The water depth contours are parallel. Initially $\theta_1 = \gamma_1$ where each initial direction indicated on the figure is the angle between the hydron velocity vector and the normal to the depth contours. Regardless of the wave period, for deep water angles of incidence between 0° and 74.8° the hydrons follow paths such that the angles increase to the depth of the geometric group speed maximum (see Figure 3), then undergo a point of inflection, and then decrease shoreward. As a hydron approaches shore its direction becomes perpendicular to the wave speed contours and the packet ray curvature approaches zero. For deep water angles of incidence equal to or greater than 74.8° the hydron trajectories turn and move parallel to shore in water of intermediate depth. As the hydron direction becomes parallel to shore the packet ray curvature tends to vanish; this is apparent in ray number 5.

For comparison, monochromatic rays are shown in Figure 2 for the same conditions considered in Figure 1. For large incident angles there is a striking difference between hydron and monochromatic trajectories. Whereas all the hydron rays do not reach shore all the monochromatic rays do. Note that the wavelet direction at each point along a hydron path in Figure 1 is the same as the direction of the corresponding monochromatic ray at the corresponding water depth.

It is interesting to compare the values of G and U when they differ due to




Figure 1. Hydron trajectories for a 20 sec wave period for waves beginning in deep water. The water depth contours are parallel, the scale of the plot is 1 cm = 4.87 km, and the sounding water depths are in meters. The initial hydron direction is shown for each ray and is the angle between the hydron velocity vector and the normal to the depth contours.




Figure 2. Monochromatic rays for comparison with the hydron trajectories in Figure 1.

refraction. As an example, for gravity water waves which begin in deep water it is found that

$$v_r = \tanh kh \quad (19)$$

$$U_r = \left(1 + \frac{2kh}{\sinh 2kh}\right) v_r \quad (20)$$

where h is the water depth. When (19) and (20) are substituted into (18) and the result is substituted into (2) it is found that

$$G = U \left[1 + \frac{(kh \sin \gamma_1 \operatorname{sech}^2 kh)^2}{1 - (\sin \gamma_1 \tanh kh)^2} \right]^{-\frac{1}{2}} \quad (21)$$

The ratio of the geometric group speed to its initial deep water value is presented for several incident angles in Figure 3. The initial hydron directions are defined as in Figure 1. The curve for $\gamma_1 = 0^\circ$ is the same as obtained for the ratio of U to the value of U in deep water. The amount by which the other curves differ from it is a measure of the difference between G and U .

Inspection of Figure 3 shows for a given γ_1 that the maximum of G/G_1 occurs at a greater value of kh than does the minimum of G/U . An increase in γ_1 causes a shift in both the minimum of G/U and the maximum of G/G_1 to larger values of kh . Further, the maximum peak tends to get flattened out. The curve for $\gamma_1 = 74.8^\circ$ is seen to stop abruptly at the maximum value of G/G_1 .

When $\gamma_1 = 30^\circ$ the maximum percentage difference of G from U is 2.70%. When $\gamma_1 = 45^\circ$ the value is 5.91%, for $\gamma_1 = 60^\circ$ it is 10.03%, and for $\gamma_1 = 74.79^\circ$ the value is 14.27%.

Reflection Points

To obtain a reflection point it is necessary that the waves propagate into deeper water and that the initial direction of the hydron ($\theta_1 = \gamma_1$) exceed a critical angle. The reflection point occurs at an intermediate water depth when, through refraction, the wavelets are turned parallel to the wave speed contours.

In Figure 4 two rays are shown in which the wave period is 20 sec and the




Figure 3. The variation of the geometric group speed with kh for hydrons beginning in deep water. The initial value of the geometric group speed is G_1 . The hydron directions are defined as in Figure 1.




Figure 4. Hydron trajectories for a 20 sec wave period for waves which begin at an intermediate water depth. The scale of the plot is 1 cm = 3.10 km and the sounding water depths are in meters. The initial hydron direction is shown for each ray and is defined as in Figure 1.

initial water depth is 15 m. For this case a reflection point occurs if $\theta_1 \geq 22.2^\circ$. Ray number 1 reaches deep water since $\theta_1 = 15^\circ$. For ray number 2, $\theta_1 = 23^\circ$, and a reflection point occurs at a water depth of 200 m. The variation of the hydron and wavelet directions with water depth for this ray are shown in Figure 5. The wavelet angle increases continuously with water depth and becomes parallel to the water depth contours at the reflection point. The hydron angle first increases, goes through a maximum, and then becomes perpendicular to the water depth contours at the reflection point. This is clearly seen in Figure 4.

Plots of how G and U vary with water depth in approaching the reflection point are shown in Figure 5. Both speeds have a maximum value. However, there is a marked contrast between G and U at the reflection point where the geometric group speed is zero.

In Figure 6 the ratio of the packet ray curvature to its initial value is sketched as a function of the wavelet direction for ray number 2 in Figure 4. The most striking feature of this curve is that within about 2° of the reflection point the packet ray curvature suddenly goes to infinity.

It is interesting to observe that the velocity of the wave packet goes to zero at the reflection point. This is exactly what the velocity of a particle should do when there is reflection. Therefore, in water waves, as in quantum mechanics, the wave-particle duality is encountered.

Summary

Refraction causes a wave packet (hydron) trajectory to become directed either parallel or perpendicular to the water depth contours. In either case the packet ray curvature will vanish. For hydrons propagating toward deep water, if the initial direction exceeds a critical angle total reflection occurs. At the reflection point the wavelet direction becomes parallel to the wave speed contours, the hydron direction becomes perpendicular to the contours, the geometric group velocity goes to zero, and the packet ray curvature becomes infinite.




Figure 5. Variations of the wavelet direction γ , hydron direction θ , the speed $U = d\omega/dk$, and the geometric group speed $G = U \cos (\theta - \gamma)$ up to the reflection point for ray number 2 in Figure 4.




Figure 6. The ratio of the hydron ray curvature to its absolute initial value as a function of the wavelet direction up to the reflection point for ray number 2 in Figure 4.

References

- Arthur, R.S., W. H. Munk, and J. D. Isaacs, The direct construction of wave rays, Trans, AGU, 33, 855-865, 1952.
- Breeding, J.E., Jr., Velocities and refraction laws of wave groups: a verification, J. Geophys. Res., in press.
- Breeding, J. E., Jr., K. C. Matson, and N. Riahi, A method for calculating wave packet trajectories and wave heights, Department of Oceanography, Florida State University, Tallahassee, 1978.
- Munk, W. H., and R. S. Arthur, Wave intensity along a refracted ray, in Gravity Waves, pp. 95-108, National Bureau of Standards Circular 521, Washington, D. C., 1952.
- Purser, W. F. C., and J. L. Synge, Water waves and Hamilton's method, Nature, 194, 268, 1962.
- Synge, J. L., Water Waves and hydrons, Science, 138, 13-15, 1962.

DISTRIBUTION LIST

Office of Naval Research 2
Geography Programs
Code 462
Arlington, VA 22217

Defense Documentation Center 12
Cameron Station
Alexandria, VA 22314

Director, Naval Research Lab 6
Attention Technical Information
Officer
Washington, DC 20375

Director
Office of Naval Research Branch
Office
536 South Clark St.
Chicago, Illinois 60605

Director
Office of Naval Research Branch
Office
495 Summer St.
Boston, MA 02210

Commanding Officer
Office of Naval Research Branch
Office
Box 39
FPO New York 09510

Chief of Naval Research
Asst. for Marine Corps Matters
Code 100M
Office of Naval Research
Arlington, VA 22217

Office of Naval Research
Code 480
National Space Technology
Laboratories
Bay St. Louis, MS 39520

Office of Naval Research
Operational Applications Division
Code 200
Arlington, VA 22217

Office of Naval Research
Scientific Liaison Officer
Scripps Institution of
Oceanography
La Jolla, CA 92093

Director, Naval Research
Laboratory
Attn: Library, Code 2628
Washington, DC 20375

ONR Scientific Liaison Group
American Embassy, Room A-407
APO San Francisco, CA 96503

Commander
Naval Oceanographic Office
Attn: Library, Code 1600
Washington, DC 20374

Naval Oceanographic Office
Code 3001
Washington, DC 20374

Chief of Naval Operations
OP 987P1
Department of the Navy
Washington, DC 20350

Oceanographer of the Navy
Hoffman II Building
200 Stovall St.
Alexandria, VA 22322

Naval Academy Library
US Naval Academy
Annapolis, MD 21402

Commanding Officer
Naval Coastal Systems
Center
Panama City, FL 32407

Librarian
Naval Intelligence
Support Center
4301 Suitland Rd.
Washington, DC 20390

Dr. William T. Fox
Dept. of Geology
Williams College
Williamstown, MA 01267

Dr. Hsiang Wang
Dept. of Civil Engineering
Dupont Hall
University of Delaware
Newark, DE 19711

Dr. John T. Kuo
Henry Krumb School of Mines
Seely W. Mudd Building
Columbia University
New York, NY 10027

Dr. Edward B. Thornton
Dept. of Oceanography
Naval Postgraduate School
Monterey, CA 93940

Prof. C.A.M. King
Dept. of Geography
University of Nottingham
Nottingham, England NG7 2RD

Dr. Douglas L. Inman
University of California A-009
Shore Processes Laboratory
La Jolla, CA 92093

Dr. Omar Shemdin
Jet Propulsion Laboratory
183-501
4800 Oak Grove Dr.
Pasadena, CA 91103

Dr. William L. Wood
Department of Geosciences
Purdue University
Lafayette, IN 47907

Dr. Alan W. Niedoroda
Director, Coastal Research
Center
University of Massachusetts
Amherst, MA 01002

Dr. John B. Southard
Dept. of Earth and Planetary
Sciences
MIT
Cambridge, MA 02139

Dr. J. Ernest Breeding, Jr.
Dept. of Oceanography
FSU
Tallahassee, FL 32306

Dr. John C. Kraft
Dept. of Geology
University of Delaware
Newark, DE 19711

Dr. Dag Nummedal
Dept. of Geology
University of South Carolina
Columbia, SC 29208

Mr. Fred Thomson
Environmental Research
Institute
P.O. Box 618
Ann Arbor, MI 48107

Dr. Thomas K. Peucker
Simon Fraser University
Dept. of Geography
Burnaby 2, B.C., Canada

Dr. Robert Dolan
Department of Environmental
Sciences
University of Virginia
Charlottesville, VA 22903

Dr. Lester A. Gerhardt
Rennselaer Polytechnic
Institute
Troy, New York 12181

Director
Office of Naval Research
Branch Office
1030 East Green Street
Pasadena, California 91101

Dr. Yoshimi Goda, Director
Wave Research Division
Port and Harbor Research Instit.
Ministry of Transportation
1-1 Nagase, 3 Chome
Yokosuka, 239 Japan

Prof. Dr. Rer. Nat. H.G.
Gierloff-Emden
Institut F. Geographie
Universitaet Muenchen
Luisenstrasse 37/III
D-800 Muenchen 2, West Germany

Dr. R. Koester
Geol-Palaeontolog. Institut
Universitaet Kiel
Olshausenstrasse 40-60
D-2300 Kiel, West Germany

Prof. Dr. Fuehrboeter
Lehrstuhl F. Hydromechanik U.
Kuestenw
Technische Hochschule Braunschweig
Beethovenstrasse 51A
D-3300 Braunschweig, West Germany

Prof. Dr. Walter Hansen
Direktor D. Instituts F.
Meereskunde
Universitaet Hamburg
Heimhuderstrasse 71
D-2000 Hamburg 13, West Germany

Prof. Dr. Klaus Hasselmann
Institute F. Geophysik
Universitaet Hamburg
Schlueterstrasse 22
D-2000 Hamburg 13, West Germany

Prof. Dr. Nils Jerlov
Institute for Physical
Oceanography
Kobenhavns Universitet
Haraldsgade 6
DK-2200 Kobenhavn, Denmark

Prof. Kiyoshi Horikawa
Dept. of Civil Engineering
University of Tokyo
7-3-1, Hongo, Bunkyo-Ku
Tokyo 113, Japan

Prof. Dr. Eugen Seibold
Geol-Palaeontolog. Institut
Universitaet Kiel
Olshausenstrasse 40-60
D-2300 Kiel, West Germany

Prof. Yuji Iwagaki
Civil Engineering Dept.
Kyoto University
9 Shimogamo Zenbuchi, Sakyo-Ku
Kyoto, Japan

Dr. H.J. Schoemaker
Waterloopkundig Laboratorium
Te Delft
61 Raam, Delft, Netherlands

Ir. M.W. Van Batenberg
Physisch Laboratorium TNO
Oude Waalsdorper Weg 63,
Den Haag, Netherlands

Prof. Toshiyuki Shigemura
Civil Engineering Dept.
National Defense Academy
1-10-20 Hashirimizu
Yokosuka 239, Japan

Mr. William T. Whelan
Telecommunication Enterprises,
Inc.
Box 88
Burtonsville, MD 20730

Dr. Benno M. Brenninkmeyer, SJ
Dept. of Geology and Geophysics
Boston College
Chestnut Hill, MA 02167

Coastal Studies Institute
Louisiana State University
Baton Rouge, LA 70803

Dr. Choule J. Sonu
Tetra Tech, Inc.
630 N. Rosemead Blvd.
Pasadena, CA 91107

Dr. Richard A. Davis, Jr.
Dept. of Geology
University of South Florida
Tampa, FL 33620

Commanding Officer
Naval Civil Engineering
Laboratory
Port Hueneme, CA 93041

Officer in Charge
Environmental Research
Production Facility
Naval Postgraduate School
Monterey, CA 93940

Director
Amphibious Warfare Board
US Atlantic Fleet
Naval Amphibious Base
Norfolk, Little Creek, VA 23520

Commander, Amphibious Force
US Pacific Fleet
Force Meteorologist
Comphibpac Code 25 5
San Diego, CA 92155

Commanding General
Marine Corps Development and
Educational Command
Quantico, VA 22134

Dr. A. L. Slafkosky
Scientific Advisor
Commandant of the Marine Corps
Code Mc-Rd-1
Washington, DC 20380

Defense Intelligence Agency
Central Reference Division
Code RDS-3
Washington, DC 20301

Director
Coastal Engineering Research
Center
US Army Corps of Engineers
Kingman Building
Fort Belvoir, VA 22060

Chief, Wave Dynamics Division
USAE-WES
PO Box 631
Vicksburg, MS 39180

Commandant
US Coast Guard
ATTN: GECV/61
Washington, DC 20591

Office of Research and
Development
DS/62
US Coast Guard
Washington, DC 20591

National Oceanographic Data
Center D764
Environmental Data Services
NOAA
Washington, DC 20235

Central Intelligence Agency
Attn: OCR/DD-Publications
Washington, DC 20505

Dr. Donald Swift
Marine Geology and Geophysics
Laboratory
15 Rickenbacker Causeway
AOML-NOAA
Miami, FL 33149

MinisterialDirektor
Dr. F. Wever
RUE/FO
Bundesministerium der Verteidigung
Hardthoehe
D-5300 Bonn, West Germany

Oberregierungsrat
Dr. Ullrich
RUE/FO
Bundesministerium der Verteidigung
Hardthoehe
D-5300 Bonn, West Germany

Mr. Tage Strarup
Defense Research Establishment
Osterbrogades Kaserne
DK-2100 Kobenhavn O, Denmark

Dr. Warren C. Thompson
Dept. of Meteorology &
Oceanography
Naval Postgraduate School
Monterey, California 93940

Block 20.

to turn the hydron so that it is directed either perpendicular or parallel to the shoreline. The first case is similar to monochromatic waves. It is the result for hydrons which begin in deep water if the angle of incidence is between 0° and 74.8° with respect to the contour normal. However, for deep water angles of incidence equal to or greater than 74.8° , the hydrons are turned and move parallel to shore in water of intermediate depth. The packet ray curvature approaches infinity as the wavelet direction, but not the hydron direction, becomes parallel to the wave speed contours. The result is total reflection of the waves. Total reflection occurs if a hydron is moving into deeper water and its initial direction exceeds a critical angle. At the reflection point the hydron direction becomes perpendicular to the water depth contours. Further, the hydron velocity goes to zero, which is consistent with a particle concept. As in quantum mechanics, the wave-particle duality is encountered.